

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
11.06.2003 Bulletin 2003/24

(51) Int Cl.7: **A61K 31/505**, A61P 21/00,
A61P 29/02, A61P 43/00

(21) Application number: **02258123.5**

(22) Date of filing: **26.11.2002**

<div>(84) Designated Contracting States: AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR Designated Extension States: AL LT LV MK RO SI</div> <div>(30) Priority: 06.12.2001 GB 0129274</div> <div>(71) Applicants:<ul style="list-style-type: none">• Pfizer Limited Sandwich, Kent CT13 9NJ (GB) Designated Contracting States: GB• PFIZER INC. New York, N.Y. 10017 (US) Designated Contracting States: AT BE BG CH CY CZ DE DK EE ES FI FR GR IE IT LU MC NL PT SE TR SK</div>	<div>(72) Inventors:<ul style="list-style-type: none">• Abel, Samantha Sandwich, Kent CT13 9NJ (GB)• Ellis, Peter Sandwich, Kent CT13 9NJ (GB)</div> <div>(74) Representative: Wood, David John et al UK Patent Department, Pfizer Limited, Ramsgate Road, Sandwich, Kent CT13 9NJ (GB)</div>
--	---

(54)

Kit for reducing aching caused by pde-v inhibitors

(57) The present invention relates to kits, and aspects thereof, for reducing or eliminating the aching associated with the administration of multiple doses of a PDE5 inhibitor, the kits comprising a plurality of pharmaceutical compositions for sequential administration

over a period of time which compositions comprise increasing amounts of PDE5 inhibitor starting with an amount which gives a sub-optimal response and ending with an amount which gives an optimal response.

EP 1 317 924 A1

Description

[0001] The present invention relates to kits, and aspects thereof, for the sequential administration of cyclic guanosine 3', 5'-monophosphate phosphodiesterase Type 5 inhibitors, by the use of which kits the aching associated with the multiple dosing of such inhibitors is eliminated or significantly reduced.

[0002] Compounds which inhibit cyclic guanosine 3', 5'-monophosphate phosphodiesterase Type 5, commonly referred to as cGMP PDE5 inhibitors or simply PDE5i's, such as those described in International Patent Application No. WO 94/28902, have been found to be potent and effective compounds for the treatment of male erectile dysfunction (MED), impotence and female sexual disorders. This finding led to the development of the compound sildenafil (5-[2-ethoxy-5-(4-methyl-1-piperazinylsulphonyl)phenyl]-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one) (Viagra™) which has proved outstandingly successful as the first orally effective treatment for MED.

[0003] Following the success of sildenafil in treating MED, PDE5 inhibitors have been shown to have efficacy in the treatment of other medical conditions, such as neuropathy (EP 1129706), diabetic ulcers (WO 01/51042) and pulmonary hypertension (EP 1097711).

[0004] While MED has been found to be effectively treated by a single dose immediately prior to sexual activity, other diseases, such as those mentioned above, typically require a prolonged course of PDE5i treatment. This can involve dosing with a PDE5 inhibitor up to 3 (tid) or 4 times a day over a period of several days.

[0005] Examples of conditions for which multiple daily dosing of a PDE5 inhibitor is indicated include the aforementioned neuropathy, diabetic ulcers (particularly diabetic foot ulcers) and pulmonary hypertension plus conditions selected from the group comprising diabetes, hypertension, angina (including vasospastic angina), heart failure, intermittent claudication; post-angioplasty stenosis; cardioprotection (ischaemic preconditioning); atherosclerosis; acute coronary syndromes; chronic obstructive pulmonary disease (COPD); corpulmonale; Eisenmenger's syndrome; Raynaud's syndrome; systemic scleroderma; pre-eclampsia; intrauterine growth retardation (IUGR); infertility; preterm labour, recurrent abortion, dysmenorrhoea, stroke, Alzheimer's disease, cognitive impairment, Parkinson's and other degenerative disorders, glaucoma, macular degeneration, myasthenia gravis, benign prostatic hyperplasia (BPH), lower urinary tract syndrome (LUTS)/overactive bladder (OAB), chronic inflammatory hyperalgesia, neuropathic pain, back pain, chronic lymphatic leukaemia (CLL)/apoptosis, insulin resistance syndrome, acne and anal fissures

[0006] While the foregoing are all non-sexual conditions, it is possible to envisage a situation wherein multiple dosing of a PDE5 inhibitor might be useful in the long-term treatment of persistent sexual conditions, for example, the aforementioned MED and female sexual disorders. The latter include female sexual dysfunction (FSD), clitoral dysfunction, female sexual arousal disorder, female sexual pain disorder or female sexual orgasmic dysfunction (FSOD), sexual dysfunction due to spinal cord injury and selective serotonin reuptake inhibitor (SSRI)-induced sexual dysfunction.

[0007] A problem associated with the multiple daily dosing of PDE5 inhibitors is that subjects experience, as a side effect, aching in certain parts of the body. Subjects describe this aching as predominantly a dull aching without muscle tenderness. Aching is experienced predominantly in the "postural muscles", typically the calves, thighs, buttocks, lower back, shoulders and neck regions. These symptoms may be described as myalgia or back pain; for the purposes of this application, they will be referred to as "aching". Often the aching is painful and debilitating and in some individuals can result in their stopping a course of treatment with a PDE5 inhibitor.

[0008] Aching as a side effect appears to be a class effect associated with the multiple dosing of any PDE5 inhibitor. With sildenafil, aching generally commences on the second or third day of a multiple dosing regime. At higher doses (for example, 75mg tid), there is evidence that aching disappears by the sixth or seventh day. At higher doses (75mg tid), the incidence of aching was 100%, falling to 60% at a dose of 25mg tid. The effect has also been observed with the PDE5 inhibitors 3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-(2-methoxyethoxy)pyridin-3-yl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-d]-pyrimidin-7-one (WO 99/54333, Example 4) and 3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-*n*-propoxyphenyl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (WO 98/49166, Example 4). The PDE5 inhibitor tadalafil (Cialis™) has also been reported to cause backache and myalgia (H Padma-Nathan *et al*, Int. J. Impotence Res., 2001, 12, 2-9 and the presentation by Eli Lilly/ICOS at the European Society for Sexual and Impotence Research meeting held in Rome between 30 September 2001 and 3 October 2001).

[0009] We have now unexpectedly found that the severity of the aching associated with multiple dosing of PDE5 inhibitors may be significantly reduced by administering the PDE5 inhibitor in a defined sequence over a period of time, specifically by starting with an amount of PDE5 inhibitor which causes a sub-optimal response and ending with an amount which causes an optimal response. The amount of PDE5 inhibitor used to give the sub-optimal response is typically from 1 to 50%, preferably from 1 to 25%, of the amount which gives the optimal response, optimal response being defined as the desired clinical response with a minimum of non-aching side effects.

[0010] This finding was particularly surprising as sequential administration has previously been applied to relieve acute side effects experienced after a first single dose. Sequential administration would not have been expected to limit side effects which were experienced after several days, since late onset side effects are due to overall total exposure to drug over time, rather than the typical first exposure side effects which are directly associated with the plasma

concentration following a high first dose.

[0011] We have also surprisingly found that when the subject is multiply dosed in accordance with the invention, if the patient misses one or more doses (or even an entire daily dose) and resumes taking the PDE5 inhibitor at the same dose, aching does not re-occur.

5 [0012] Sequential administration in accordance with the invention of a PDE5 inhibitor selected from sildenafil, 3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-(2-methoxyethoxy)pyridin-3-yl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (WO 99/54333, Example 4), 3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-*n*-propoxyphenyl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (WO 98/49166, Example 4) and 2-[2-ethoxy-5-(4-ethyl-piperazin-1-yl-1-sulphonyl)-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-f][1,2,4]triazin-4-one (vardenafil) has
10 indicated that for the treatment of neuropathy, diabetic ulcers (particularly diabetic foot ulcers), pulmonary hypertension and diabetes the amount of PDE5 inhibitor giving a sub-optimal response is typically in the range of from 1 to 25mg and the amount which causes an optimal response is in the range of from 25 to 100mg.

[0013] According to a first aspect of the invention, there is provided a kit for reducing the aching associated with the administration of multiple doses of a PDE5 inhibitor, said kit comprising a plurality of pharmaceutical compositions for
15 sequential administration over a period of time which compositions comprise increasing amounts of PDE5 inhibitor starting with an amount which gives a sub-optimal response and ending with an amount which gives an optimal response.

[0014] According to three preferred embodiments of the first aspect of invention, said kit ideally comprises

20 (i) from 2 to 6 PDE5i compositions for sequential administration over 2 days wherein the amount in those compositions which give a sub-optimal response comprises from 1 to 50%, preferably from 1 to 25%, of the amount which gives an optimal response;

25 (ii) from 14 to 42 PDE5i compositions for sequential administration over 14 days wherein the amount in those compositions which give a sub-optimal response comprises from 1 to 50%, preferably from 1 to 25%, of the amount which gives an optimal response; and

30 (iii) compositions comprising (a) from 1 to 15mg sildenafil for administration from 1 to 3 times a day for 3 to 14 days and (b) compositions comprising from 25 to 100mg of sildenafil for administration for the remainder of the prescribed treatment period.

[0015] By "prescribed treatment period" is meant the period that the physician deems it necessary to administer the drug at therapeutically effective doses.

35 [0016] According to another aspect of the invention, there is provided the use of a PDE5 inhibitor in the manufacture of a pharmaceutical composition for the treatment of any condition for which a PDE5 inhibitor is indicated, which composition is one of the plurality of pharmaceutical compositions used in the kit of the invention.

[0017] Preferred embodiments of this use include the manufacture of pharmaceutical compositions in accordance with those found in the preferred embodiments of the kit (*supra*) for the treatment of one or more of the aforementioned conditions.

40 [0018] According to a further aspect of the invention, there is provided a method for the treatment of any condition for which multiple dosing of a PDE5 inhibitor is indicated, which method comprises the sequential administration of a plurality of pharmaceutical compositions over a period of time which compositions are in accordance with those used in the kit of the invention.

45 [0019] Preferred embodiments of this method of treatment include those methods which employ a plurality of compositions in accordance with those found in the preferred embodiments of the kit (*supra*) in the treatment of one or more of the aforementioned conditions.

[0020] In all of the aforementioned aspects of the invention, the PDE5 inhibitor which gives the sub-optimal response may be different from the PDE5 inhibitor which gives the optimal response.

[0021] PDE5 inhibitors suitable for use in all aspects of the invention include

50 pyrazolo[4,3-d]pyrimidin-7-ones as disclosed in EP 0463756;

pyrazolo[4,3-d]pyrimidin-7-ones as disclosed in EP 0526004;

55 pyrazolo[4,3-d]pyrimidin-7-ones as disclosed in WO 93/06104;

isomeric pyrazolo[3,4-d]pyrimidin-4-ones as disclosed in WO 93/07149;

quinazolin-4-ones as disclosed in WO 93/12095;

pyrido[3,2-d]pyrimidin-4-ones as disclosed in WO 94/05661;

5 purin-6-ones as disclosed in WO 94/00453;

pyrazolo[4,3-d]pyrimidin-7-ones as disclosed in WO 98/49166;

10 pyrazolo[4,3-d]pyrimidin-7-ones as disclosed in WO 99/54333;

pyrazolo[4,3-d]pyrimidin-4-ones as disclosed in EP 0995751;

pyrazolo[4,3-d]pyrimidin-7-ones as disclosed in WO 00/24745;

15 pyrazolo[4,3-d]pyrimidin-4-ones as disclosed in EP 0995750;

compounds disclosed in WO 95/19978;

20 compounds disclosed in WO 99/24433;

compounds disclosed in WO 93/07124;

pyrazolo[4,3-d]pyrimidin-7-ones as disclosed in WO 01/27112;

25 pyrazolo[4,3-d]pyrimidin-7-ones as disclosed in WO 01/27113;

compounds disclosed in EP 1092718; and

compounds disclosed in EP 1092719.

30

[0022] Specific PDE5 inhibitors suitable for use in all aspects of the invention include

4-bromo-5-(pyridylmethylamino)-6-[3-(4-chlorophenyl)-propoxy]-3(2H)pyridazinone;

35

1-[4-[(1,3-benzodioxol-5-ylmethyl)amino]-6-chloro-2-quinazolinyl]-4-piperidine-carboxylic acid, monosodium salt;

(+)-*cis*-5,6a,7,9,9a-hexahydro-2-[4-(trifluoromethyl)-phenylmethyl-5-methyl-cyclopent-4,5]imidazo[2,1-b]purin-4(3H)one;

40

furazlocillin;

cis-2-hexyl-5-methyl-3,4,5,6a,7,8,9,9a-octahydrocyclopent[4,5]-imidazo[2,1-b]purin-4-one;

3-acetyl-1-(2-chlorobenzyl)-2-propylindole-6-carboxylate;

45

3-acetyl-1-(2-chlorobenzyl)-2-propylindole-6-carboxylate;

4-bromo-5-(3-pyridylmethylamino)-6-(3-(4-chlorophenyl)propoxy)-3-(2H)pyridazinone;

50

l-methyl-5(5-morpholinoacetyl-2-n-propoxyphenyl)-3-n-propyl-1,6-dihydro-7H-pyrazolo(4,3-d)pyrimidin-7-one;

1-[4-[(1,3-benzodioxol-5-ylmethyl)amino]-6-chloro-2-quinazolinyl]-4-piperidinecarboxylic acid, monosodium salt;

compounds of Pharmaprojects No. 4516 (Glaxo Wellcome);

55

compounds of Pharmaprojects No. 5051 (Bayer);

compounds of Pharmaprojects No. 5064 (Kyowa Hakko; WO 96/26940);

compounds of Pharmaprojects No. 5069 (Schering Plough);

GF-196960 (Glaxo Wellcome);

5 E-8010 and E-4010 (Eisai);

Bay-38-3045 & 38-9456 (Bayer); and

Sch-51866 (Schering Plough).

10

[0023] Particularly preferred PDE5 inhibitors suitable for use in all aspects of the invention include

15

5-[2-ethoxy-5-(4-methyl-1-piperazinylsulphonyl)phenyl]-1-methyl-3-n-propyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (sildenafil), also known as 1-[[3-(6,7-dihydro-1-methyl-7-oxo-3-propyl-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-4-ethoxyphenyl]sulphonyl]-4-methylpiperazine (EP 0463756);

5-(2-ethoxy-5-morpholinoacetylphenyl)-1-methyl-3-n-propyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (EP 0526004);

20

(6R,12aR)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione (WO 95/19978, Examples 1, 3, 7, 8, 78 and 95) (tadalafil);

25

2-[2-ethoxy-5-(4-ethylpiperazin-1-yl)-1-sulphonyl]-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-f][1,2,4]triazin-4-one, also known as 1-[[3-(3,4-dihydro-5-methyl-4-oxo-7-propylimidazo[5,1-f]-as-triazin-2-yl)-4-ethoxyphenyl]sulphonyl]-4-ethylpiperazine (WO99/24433, Examples 20, 19, 336 and 337) (vardenafil);

30

5-[2-ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-yl]-3-ethyl-2-[2-methoxyethyl]-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one, also known as 1-{6-ethoxy-5-[3-ethyl-6,7-dihydro-2-(2-methoxyethyl)-7-oxo-2H-pyrazolo[4,3-d]pyrimidin-5-yl]-3-pyridylsulphonyl}-4-ethylpiperazine (WO 01/27113, Example 8);

5-(5-Acetyl-2-butoxy-3-pyridinyl)-3-ethyl-2-(1-ethyl-3-azetidiny)-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (WO 01/27112, Example 132);

35

3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-(2-methoxyethoxy)pyridin-3-yl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (WO 99/54333, Example 4);

3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-n-propoxyphenyl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (WO 98/49166, Example 4);

40

(+)-3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-(2-methoxy-1(R)-methylethoxy)pyridin-3-yl]-2-methyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one, also known as 3-ethyl-5-{5-[4-ethylpiperazin-1-ylsulphonyl]-2-[(1 R)-2-methoxy-1-methylethyl]oxy}pyridin-3-yl]-2-methyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (WO 99/54333);

45

5-[2-*iso*-Butoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-yl]-3-ethyl-2-(1-methylpiperidin-4-yl)-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (WO 01/27113, Example 15);

50

5-[2-Ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-yl]-3-ethyl-2-phenyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (WO 01/27113, Example 66);

5-(5-Acetyl-2-propoxy-3-pyridinyl)-3-ethyl-2-(1-isopropyl-3-azetidiny)-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (WO 01/27112, Example 124); the compound of Example 11 of International Patent Application No. WO 93/07124 (Eisai); and

55

compounds 3 and 14 of Rotella D P, *J Med Chem*, 43, 1257 (2000).

[0024] The contents of the published patent applications and journal articles, particularly the general formulae of the therapeutically active compounds of the claims and compounds exemplified therein, are incorporated herein by refer-

ence.

[0025] Of the aforementioned compounds,

5 5-[2-ethoxy-5-(4-methyl-1-piperazinylsulphonyl)phenyl]-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one (sildenafil);

5-(2-ethoxy-5-morpholinoacetylphenyl)-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one;

10 (6*R*,12*aR*)-2,3,6,7,12,12*a*-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-*b*]indole-1,4-dione (tadalafil);

2-[2-ethoxy-5-(4-ethyl-piperazin-1-yl-1-sulphonyl)-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-*f*][1,2,4]triazin-4-one (vardenafil);

15 5-[2-ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-yl]-3-ethyl-2-[2-methoxyethyl]-2,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one;

5-(5-acetyl-2-butoxy-3-pyridinyl)-3-ethyl-2-(1-ethyl-3-azetidiny)-2,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one;

20 3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-(2-methoxyethoxy)pyridin-3-yl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one;

3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-*n*-propoxyphenyl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one);

25 and pharmaceutically acceptable salts thereof are particularly preferred.

[0026] And of these compounds,

30 5-[2-ethoxy-5-(4-methyl-1-piperazinylsulphonyl)phenyl]-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one (sildenafil);

5-(2-ethoxy-5-morpholinoacetylphenyl)-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one;

and pharmaceutically acceptable salts thereof are especially preferred, particularly the citrate of the former.

35 [0027] The following Examples derived from studies involving the sequential administration of the PDE5 inhibitors sildenafil and 5-(2-ethoxy-5-morpholinoacetylphenyl)-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one (Compound A) serve to illustrate the invention.

40 EXAMPLES

[0028] All adverse events observed or recorded during the studies were recorded by the investigator, including information about concomitant illnesses and therapeutic interventions. For all adverse events, the investigator pursued and obtained information adequate to determine the outcome of the adverse event. Physical examinations at screening and medical histories of all subjects were normal with no indication of any joint, muscle, or tendon weakness.

45 [0029] The Examples typically commence at a low dose of PDE5 inhibitor, for example, 2.5mg three times a day (tid) for Compound A (Examples 4, 6 and 7) with periodic dose escalation aimed at achieving a final regimen of 20mg tid. By means of such dose escalation, it was possible to significantly reduce the aching previously associated with the multiple dosing of PDE5 inhibitors.

50 STUDY 1 - SILDENAFIL

Example 1

(Placebo and comparative - negative and positive controls)

55 [0030] In a double blind, placebo-controlled, multiple dose study, from 8 to 10 subjects were given 5-[2-ethoxy-5-(4-methyl-1-piperazinylsulphonyl)phenyl]-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one (sildenafil) three times a day at one of three different doses (25mg tid, 50mg tid, or 75mg tid). Dosage took the form

of oral capsules.

[0031] The most common adverse events were back pain, headache and myalgia.

[0032] The results for myalgia and back pain ("aching") were as follows:

TABLE 1

Dosage	Placebo	25mg tid	50mg tid	75mg tid
Myalgia	0/8	2/10	5/9	3/8
Back pain	0/8	2/10	2/9	5/8

[0033] At the lower doses (25mg tid and 50mg tid), headache and myalgia typically occurred after a few days of treatment, but did not persist with further treatment.

[0034] At the higher dose (75mg tid), there were three reports of severe treatment-related adverse events which led to discontinuance of the treatment for the subjects in question. These subjects experienced respectively (1) severe myalgia (lower back pain) on Day 27 after 5 days of tid dosing, treatment discontinued on Day 30, (2) severe lower back pain on Day 27 after 4 days of tid dosing, treatment discontinued on Day 31 and (3) severe pain (throbbing pain behind thighs) on Day 23 after 2 days of tid dosing, treatment discontinued on Day 24.

STUDY 2 - COMPOUND A

Example 2

(Placebo - negative control)

[0035] Nine subjects were given placebo three times a day (tid) for 21 days.

[0036] None of the subjects reported any myalgia or back pain ("aching"):

TABLE 2

	Mild/moderate	Severe
Myalgia/back pain	0/9	0/9

Example 3

(Comparative - positive control)

[0037] Six subjects were given 20mg of Compound A three times a day (tid) for 7 days. Dosage took the form of immediate release tablets.

[0038] The results for myalgia and back pain ("aching") were as follows:

TABLE 3

	Mild/moderate	Severe
Myalgia/back pain	0/6	6/6

[0039] Due to the severity of the aching, four subjects had to be withdrawn after only 4 days of treatment.

Example 4

[0040] Nine subjects were given 2.5mg of Compound A three times a day (tid) for 7 days, then 7.5mg tid for 7 days, then 20mg tid for a further 7 days. Dosage took the form of immediate release tablets.

[0041] The results for myalgia and back pain ("aching") were as follows:

TABLE 4

	Mild/moderate	Severe
Myalgia/back pain	1/9	1/9

CONCLUSION

[0042] It can be seen from Table 4 that those subjects who followed the escalating dosing regime of Example 4 experienced a significant reduction in aching and had no withdrawals compared with those subjects who followed the non-escalating dosing regime of (comparative) Example 3.

STUDY 3 - COMPOUND A

Example 5

(Placebo - negative control)

[0043] Nine subjects were given placebo three times a day (tid) for 21 days.

[0044] The results for myalgia and back pain ("aching") were as follows:

TABLE 5

	Mild	Moderate	Severe
Myalgia	4/9	0/9	0/9
Back pain	0/9	0/9	0/9

Example 6

[0045] Nine subjects were given 2.5mg of Compound A three times a day (tid) for 7 days followed by 20mg tid for a further 7 days. The subjects were given a 24-hour drug holiday, *i.e.* a day without taking the inhibitor, then 20mg tid dosing was continued for another 6 days. Dosage took the form of immediate release tablets.

[0046] The results for myalgia and back pain ("aching") were as follows:

TABLE 6

	Mild	Moderate	Severe
Myalgia	2/9	3/9	1/9
Back pain	2/9	0/9	0/9

[0047] There was no apparent increase in the incidence of aching after the 24-hour drug-free period.

Example 7

[0048] Nine subjects were given 2.5mg of Compound A three times a day (tid) for 4 days, then 7.5mg tid for 3 days, then 20mg tid for a further 7 days. The subjects were given a 24-hour drug holiday, *i.e.* a day without taking the inhibitor, then 20mg tid dosing was continued for another 6 days. Dosage took the form of immediate release tablets.

[0049] The results for myalgia and back pain ("aching") were as follows:

TABLE 7

	Mild	Moderate	Severe
Myalgia	4/9	1/9	0/9
Back pain	0/9	0/9	0/9

[0050] There was no apparent increase in the incidence of aching after the 24-hour drug-free period.

CONCLUSION

[0051] It can be seen from Tables 6 and 7 that those subjects who followed the escalating dosing regimes of Examples 6 and 7 experienced a significant reduction in aching and had no withdrawals compared with those subjects who followed the non-escalating dosing regime of (comparative) Example 3. Triple dosing (Example 7) appeared to be particularly effective in reducing aching.

[0052] It was also observed that the 24-hour drug-free period of Examples 6 and 7 did not lead to an increased incidence of aching when 20mg tid dosing was resumed. Thus strict compliance with the escalating dosing regime does not appear to be necessary in order to achieve the desired effect.

5

Claims

1. A kit for reducing the aching associated with the administration of multiple doses of a PDE5 inhibitor, said kit comprising a plurality of pharmaceutical compositions for sequential administration over a period of time which compositions comprise increasing amounts of PDE5 inhibitor starting with an amount which gives a sub-optimal response and ending with an amount which gives an optimal response.
2. A kit according to Claim 1, wherein said kit comprises from 2 to 6 pharmaceutical compositions for sequential administration over 2 days which compositions comprise increasing amounts of PDE5 inhibitor starting with from 1 to 50% of the amount which gives an optimal response and ending with an amount which gives an optimal response.
3. A kit according to Claim 1, wherein said kit comprises from 14 to 42 pharmaceutical compositions for sequential administration over 14 days which compositions comprise increasing amounts of PDE5 inhibitor starting with from 1 to 50% of the amount which gives an optimal response and ending with an amount which gives an optimal response.
4. A kit according to Claim 2 or 3, wherein said compositions comprise increasing amounts of PDE5 inhibitor starting with from 1 to 25% of the amount which gives an optimal response and ending with an amount which gives an optimal response.
5. A kit according to any of Claims 1 to 4, wherein the PDE5 inhibitor which gives the sub-optimal response is different from the PDE5 inhibitor which gives the optimal response.
6. A kit according to any of Claims 1 to 5 wherein the PDE5 inhibitor(s) is one or more selected from the group comprising
 - 5-[2-ethoxy-5-(4-methyl-1-piperazinylsulphonyl)phenyl]-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one (sildenafil);
 - 5-(2-ethoxy-5-morpholinoacetylphenyl)-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one;
 - (6R,12aR)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-*b*]indole-1,4-dione (tadalafil);
 - 2-[2-ethoxy-5-(4-ethyl-piperazin-1-yl-1-sulphonyl)-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-*f*][1,2,4]triazin-4-one (vardenafil);
 - 5-[2-ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-yl]-3-ethyl-2-[2-methoxyethyl]-2,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one;
 - 5-(5-acetyl-2-butoxy-3-pyridinyl)-3-ethyl-2-(1-ethyl-3-azetidiny)-2,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one;
 - 3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-*n*-propoxyphenyl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one;
 - 3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-(2-methoxyethoxy)pyridin-3-yl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one;
 and pharmaceutically acceptable salts thereof.
7. A kit according to Claim 6 wherein the PDE5 inhibitor is

5-[2-ethoxy-5-(4-methyl-1-piperazinylsulphonyl)phenyl]-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one (sildenafil);

5-(2-ethoxy-5-morpholinoacetylphenyl)-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one;

5

or a pharmaceutically acceptable salt of either thereof.

8. A kit for reducing the aching associated with the administration of multiple doses of sildenafil, said kit comprising a plurality of pharmaceutical compositions for sequential administration over 7 or more days which compositions comprise (a) from 1 to 15mg of sildenafil to be administered from 1 to 3 times a day for 3 to 14 days and (b) from 25 to 100mg of sildenafil to be administered for the remainder of the prescribed treatment period.

10

9. The use of a PDE5 inhibitor in the manufacture of a pharmaceutical composition for the treatment of any condition for which multiple dosing of said inhibitor is indicated, wherein said composition is one of a plurality of pharmaceutical compositions for sequential administration over a period of time which compositions comprise increasing amounts of PDE5 inhibitor starting from an amount which gives a sub-optimal response and ending with an amount which gives an optimal response.

15

10. Use according to Claim 9, wherein said composition comprises one of from 2 to 6 pharmaceutical compositions for sequential administration over a period of 2 days which compositions comprise increasing amounts of PDE5 inhibitor starting with an amount comprising from 1 to 50% of the amount which gives an optimal response and ending with an amount which gives an optimal response.

20

11. Use according to Claim 9, wherein said composition comprises one of from 14 to 42 pharmaceutical compositions for sequential administration over a period of 14 days which compositions comprise increasing amounts of PDE5 inhibitor starting with an amount comprising from 1 to 50% of the amount which gives an optimal response and ending with an amount which gives an optimal response.

25

12. Use according to Claim 10 or 11, wherein said compositions comprise increasing amounts of PDE5 inhibitor starting with an amount comprising from 1 to 25% of the amount which gives an optimal response and ending with an amount which gives an optimal response.

30

13. Use according to any of claims 9 to 12, wherein the PDE5 inhibitor which gives the sub-optimal response is different from the PDE5 inhibitor which gives the optimal response.

35

14. Use according to any of Claims 9 to 13, wherein the PDE5 inhibitor(s) is one or more selected from the group comprising

40

5-[2-ethoxy-5-(4-methyl-1-piperazinylsulphonyl)phenyl]-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one (sildenafil);

5-(2-ethoxy-5-morpholinoacetylphenyl)-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one;

45

(6*R*,12*aR*)-2,3,6,7,12,12*a*-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-*b*]indole-1,4-dione (tadalafil);

2-[2-ethoxy-5-(4-ethylpiperazin-1-yl-1-sulphonyl)-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-*f*][1,2,4]triazin-4-one (vardenafil);

50

5-[2-ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-yl]-3-ethyl-2-[2-methoxyethyl]-2,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one;

5-(5-acetyl-2-butoxy-3-pyridinyl)-3-ethyl-2-(1-ethyl-3-azetidiny)-2,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one;

55

3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-*n*-propoxyphenyl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-*d*]pyrimidin-7-one;

3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-(2-methoxyethoxy)pyridin-3-yl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one;

and pharmaceutically acceptable salts thereof.

15. Use according to Claim 14 wherein the PDE5 inhibitor is

5-[2-ethoxy-5-(4-methyl-1-piperazinylsulphonyl)phenyl]-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (sildenafil);

5-(2-ethoxy-5-morpholinoacetylphenyl)-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one;

or a pharmaceutically acceptable salt of either thereof.

16. The use of sildenafil in the manufacture of a pharmaceutical composition for the treatment of any condition for which multiple dosing of said compound is indicated, wherein said composition is one of a plurality of pharmaceutical compositions for sequential administration over a period of 7 or more days and comprises either (a) from 1 to 15mg of sildenafil to be administered from 1 to 3 times a day for a period of from 3 to 14 days or (b) from 25 to 100mg of sildenafil to be administered for the remainder of the prescribed treatment period.

17. Use according to any of Claims 9 to 16, wherein said condition is selected from the group comprising neuropathy, diabetic ulcers, (particularly diabetic foot ulcers), pulmonary hypertension, diabetes, hypertension, angina (including vasospastic angina), heart failure, intermittent claudication, post-angioplasty stenosis, cardioprotection (ischaemic preconditioning), atherosclerosis, acute coronary syndromes, chronic obstructive pulmonary disease; corpulmonale; Eisenmenger's syndrome; Raynaud's syndrome, systemic scleroderma, pre-eclampsia, intrauterine growth retardation, infertility, preterm labour, recurrent abortion, dysmenorrhoea, stroke, Alzheimer's disease, cognitive impairment, Parkinson's and other degenerative disorders, glaucoma, macular degeneration, myasthenia gravis, benign prostatic hyperplasia, lower urinary tract syndrome/overactive bladder, chronic inflammatory hyperalgesia, neuropathic pain, back pain, chronic lymphatic leukaemia/apoptosis, insulin resistance syndrome, acne and anal fissures.

18. A method for the treatment of any condition for which multiple dosing of a PDE5 inhibitor is indicated, which method comprises the sequential administration of a plurality of pharmaceutical compositions over a period of time which compositions comprise increasing amounts of PDE5 inhibitor starting with an amount which gives a sub-optimal response and ending with an amount which gives an optimal response.

19. A method according to Claim 18, which method comprises the sequential administration of from 2 to 6 pharmaceutical compositions over 2 days which compositions comprise increasing amounts of PDE5 inhibitor starting with an amount comprising from 1 to 50% of the amount which gives an optimal response and ending with an amount which gives an optimal response.

20. A method according to Claim 18, which method comprises the sequential administration of from 14 to 42 pharmaceutical compositions over 14 days which compositions comprise increasing amounts of PDE5 inhibitor starting with an amount comprising from 1 to 50% of the amount which gives an optimal response and ending with an amount which gives an optimal response.

21. A method according to Claim 19 or 20, wherein said compositions comprise increasing amounts of PDE5 inhibitor starting with an amount comprising from 1 to 25% of the amount which gives an optimal response and ending with an amount which gives an optimal response.

22. A method according to any of Claims 18 to 21, wherein the PDE5 inhibitor which give the sub-optimal response is different from the PDE5 inhibitor which gives the optimal response.

23. A method according to any of Claims 18 to 22, wherein the PDE5 inhibitor(s) is one or more selected from the group comprising

5-[2-ethoxy-5-(4-methyl-1-piperazinylsulphonyl)phenyl]-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one (sildenafil);

5-(2-ethoxy-5-morpholinoacetylphenyl)-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one;

(6R,12aR)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido
[3,4-b]indole-1,4-dione (tadalafil);

2-[2-ethoxy-5-(4-ethyl-piperazin-1-yl-1-sulphonyl)-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-f][1,2,4]triazin-
4-one (vardenafil);

5-[2-ethoxy-5-(4-ethylpiperazin-1-ylsulphonyl)pyridin-3-yl]-3-ethyl-2-[2-methoxyethyl]-2,6-dihydro-7H-pyra-
zolo[4,3-d]pyrimidin-7-one;

5-(5-acetyl-2-butoxy-3-pyridinyl)-3-ethyl-2-(1-ethyl-3-azetidiny)-2,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-
7-one;

3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-*n*-propoxyphenyl]-2-(pyridin-2-yl)methyl-2,6-dihydro-7H-pyra-
zolo[4,3-d]pyrimidin-7-one);

3-ethyl-5-[5-(4-ethylpiperazin-1-ylsulphonyl)-2-(2-methoxyethoxy)pyridin-3-yl]-2-(pyridin-2-yl)methyl-2,6-di-
hydro-7H-pyrazolo[4,3-d]pyrimidin-7-one;

and pharmaceutically acceptable salts thereof.

24. A method according to Claim 23 wherein the PDE5 inhibitor is

5-[2-ethoxy-5-(4-methyl-1-piperazinylsulphonyl)phenyl]-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-d]
pyrimidin-7-one (sildenafil);

5-(2-ethoxy-5-morpholinoacetylphenyl)-1-methyl-3-*n*-propyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one;

or a pharmaceutically acceptable salt of either thereof.

25. A method for the treatment of any condition for which multiple dosing of sildenafil is indicated, which method
comprises the administration over a period of seven or more days of (a) from 1 to 15mg of sildenafil from 1 to 3
times a day for 3 to 14 days and (b) the administration of from 25 to 100mg of sildenafil for the remainder of the
prescribed treatment period.

26. A method according to any of claims 18 to 25, wherein said condition is selected from the group comprising neu-
ropathy, diabetic ulcers, (particularly diabetic foot ulcers), pulmonary hypertension, diabetes, hypertension, angina
(including vasospastic angina), heart failure, intermittent claudication, post-angioplasty stenosis, cardioprotection
(ischaemic preconditioning), atherosclerosis, acute coronary syndromes, chronic obstructive pulmonary disease;
cor pulmonale; Eisenmenger's syndrome; Raynaud's syndrome, systemic scleroderma, pre-eclampsia, intrauterine
growth retardation, infertility, preterm labour, recurrent abortion, dysmenorrhoea, stroke, Alzheimer's disease, cog-
nitive impairment, Parkinson's and other degenerative disorders, glaucoma, macular degeneration, myasthenia
gravis, benign prostatic hyperplasia, lower urinary tract syndrome/overactive bladder, chronic inflammatory hyper-
algesia, neuropathic pain, back pain, chronic lymphatic leukaemia/apoptosis, insulin resistance syndrome, acne
and anal fissures.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 02 25 8123 shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	WO 00 66099 A (LILLY ICOS LLC ;PULLMAN WILLIAM ERNEST (US); WHITAKER JOHN STEVEN) 9 November 2000 (2000-11-09) * abstract * * page 31, line 21 - page 33, line 2 *	8,16,25	A61K31/505 A61P21/00 A61P29/02 A61P43/00
D,X	WO 99 54333 A (BUNNAGE MARK EDWARD ;MATHIAS JOHN PAUL (GB); STREET STEPHEN DEREK) 28 October 1999 (1999-10-28) * abstract * * page 19, line 25 - page 20, line 14 *	8,16,25	
D,X	WO 98 49166 A (BUNNAGE MARK EDWARD ;MATHIAS JOHN PAUL (GB); STREET STEPHEN DEREK) 5 November 1998 (1998-11-05) * abstract * * page 19, line 21 - page 20, line 13 *	8,16,25	
D,X	EP 1 129 706 A (PFIZER LTD ;PFIZER (US)) 5 September 2001 (2001-09-05) * abstract * * page 6, line 9 - line 20 *	8,16,25	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			A61K
INCOMPLETE SEARCH <p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search: see sheet C</p>			
Place of search MUNICH		Date of completion of the search 18 February 2003	Examiner Taylor, G.M.
CATEGORY OF CITED DOCUMENTS <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.02 (P04C07)



European Patent
Office

INCOMPLETE SEARCH
SHEET C

Application Number
EP 02 25 8123

Although claims 16 and 25 are directed to a method of treatment of the human/animal body (Article 52(4) EPC), the search has been carried out and based on the alleged effects of the compound/composition.

Claim(s) searched incompletely:
8,16,25

Claim(s) not searched:
1-7,9-15,17-24,26

Reason for the limitation of the search:

Present claims 1-7, 9-15, 17(part), 18-24, 26(part) relate to products/ methods/ dosing regimes defined by reference to desirable characteristics or properties, namely

"with an amount which gives a sub-optimal response and ending with an amount which gives an optimal response".

The claims cover all products/ methods/ dosing regimes having this characteristic or property, whereas the application provides support within the meaning of Art. 84 EPC and/or disclosure within the meaning of Art. 83 EPC for only a very limited number of such products/methods/dosing regimes. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

Independent of the above reasoning, the claims also lack clarity (Art. 84 EPC). An attempt is made to define the products/methods/dosing regimes by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible.

In addition to the above, the following expressions are considered vague and indefinite within the meaning of Art. 84 EPC:

"a sub-optimal response";
"an optimal response".

Furthermore, present claims 1-7,9-15,17(part),18-24,26(part) relate to an extremely large number of possible products/methods/dosing regimes/medical indications. Support within the meaning of Art. 84 EPC and/or disclosure within the meaning of Art. 83 EPC is to be found, however, for the use of only one compounds (sildenafil) against one "indication" (back pain). In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the products/ methods/ dosing regimes as defined in



European Patent
Office

INCOMPLETE SEARCH
SHEET C

Application Number
EP 02 25 8123

claims 8/16/25 for the use of sildenafil dosing regimes in the reduction of back pain.

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 02 25 8123

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
D,X	WO 01 51042 A (DAVIES MICHAEL JOHN ; PFIZER LTD (GB); PFIZER (US); WOOD RALPH E (U) 19 July 2001 (2001-07-19) * abstract * * page 6, line 20 - page 7, line 5 * -----	8,16,25	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 02 25 8123

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

18-02-2003

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0066099	A	09-11-2000	AU 3012900 A	02-11-2000
			AU 4490800 A	17-11-2000
			BE 1012957 A5	05-06-2001
			BR 0003046 A	23-07-2002
			CA 2307101 A1	30-10-2000
			CH 692478 A5	15-07-2002
			CN 1292264 A	25-04-2001
			CN 1384746 T	11-12-2002
			CZ 20013879 A3	17-04-2002
			DE 10021266 A1	16-11-2000
			DE 20007861 U1	24-08-2000
			DK 200000677 A	31-10-2000
			EP 1173181 A2	23-01-2002
			FI 20000976 A	30-10-2000
			FR 2795646 A1	05-01-2001
			GB 2351663 A	10-01-2001
			GR 2000100153 A	29-12-2000
			HR 20000243 A1	31-12-2001
			HU 0001632 A2	28-05-2001
			HU 0200912 A2	28-08-2002
			IT MI20000922 A1	26-10-2001
			JP 2000336043 A	05-12-2000
			JP 2002543116 T	17-12-2002
			LT 2000035 A , B	27-11-2000
			LU 90569 A2	27-02-2002
			LV 12560 A	20-11-2000
			LV 12560 B	20-04-2001
			NL 1015027 C2	14-02-2001
			NL 1015027 A1	31-10-2000
			NO 20002097 A	26-10-2001
			NO 20015275 A	06-12-2001
			NZ 504163 A	30-11-2001
			PL 339897 A1	06-11-2000
			PT 102457 A	30-11-2000
			SE 0001518 A	31-10-2000
			SI 20361 A	30-04-2001
			SK 15632001 A3	05-02-2002
			TR 200001132 A2	22-01-2001
			WO 0066099 A2	09-11-2000
			US 6451807 B1	17-09-2002
			US 2001053780 A1	20-12-2001
			ZA 200002058 A	02-11-2000
WO 9954333	A	28-10-1999	AP 1022 A	14-11-2001
			AU 748352 B2	06-06-2002
			AU 2742599 A	08-11-1999

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 02 25 8123

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

18-02-2003

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9954333 A		BG 104949 A	29-06-2001
		BR 9909808 A	26-12-2000
		CA 2329077 A1	28-10-1999
		CN 1305478 T	25-07-2001
		EP 1073658 A1	07-02-2001
		HR 20000712 A1	30-06-2001
		HU 0102543 A2	28-11-2001
		WO 9954333 A1	28-10-1999
		JP 2002512248 T	23-04-2002
		NO 20005255 A	19-12-2000
		PL 343794 A1	10-09-2001
		SK 15532000 A3	03-12-2001
		TR 200003039 T2	22-01-2001
		US 6251904 B1	26-06-2001
		US 2001039271 A1	08-11-2001
		ZA 9902793 A	19-10-2000
WO 9849166 A	05-11-1998	AP 1002 A	14-08-2001
		AU 730670 B2	08-03-2001
		AU 7644598 A	24-11-1998
		BG 103828 A	30-06-2000
		BR 9810233 A	17-10-2000
		CN 1253561 T	17-05-2000
		EA 2057 B1	24-12-2001
		WO 9849166 A1	05-11-1998
		EP 0977756 A1	09-02-2000
		HR 980222 A1	28-02-1999
		JP 2000510485 T	15-08-2000
		NO 995211 A	25-10-1999
		NZ 338075 A	27-10-2000
		PL 336586 A1	03-07-2000
		SK 144699 A3	12-03-2001
		TR 9902646 T2	22-05-2000
		ZA 9803478 A	25-10-1999
		HU 0001389 A2	28-09-2000
EP 1129706 A	05-09-2001	AU 6665000 A	26-04-2001
		EP 1129706 A2	05-09-2001
		HU 0004120 A2	28-11-2002
		JP 2001122803 A	08-05-2001
		NZ 507690 A	25-10-2002
		ZA 200005825 A	19-04-2002
WO 0151042 A	19-07-2001	AU 2540301 A	24-07-2001
		EP 1248623 A2	16-10-2002
		WO 0151042 A2	19-07-2001

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82